

Return-to-School Strategy¹

The following is an outline of the *Return-to-School Strategy* that should be used to help student-athletes, parents, and teachers to partner in allowing the athlete to make a gradual return to school activities.

Depending on the severity and type of the symptoms present, student-athletes will progress through the following stages at different rates. If the student-athlete experiences new symptoms or worsening symptoms at any stage, they should go back to the previous stage.

Stage	Aim	Activity	Goal of each step
1	Daily activities at home that do not give the student-athlete symptoms	Typical activities during the day as long as they do not increase symptoms (i.e. reading, texting, screen time). Start at 5-15 minutes at a time and gradually build up.	Gradual return to typical activities.
2	School activities	Homework, reading or other cognitive activities outside of the classroom.	Increase tolerance to cognitive work.
3	Return to school part- time	Gradual introduction of schoolwork. May need to start with a partial school day or with increased breaks during the day.	Increase academic activities.
4	Return to school full- time	Gradually progress.	Return to full academic activities and catch up on missed school work.

4. Sport-Specific Return-to-Sport Strategy¹

Stage	Aim	Activity	Goal of each step
1	Symptom- limiting activity	Daily activities that do not provoke symptoms.	Gradual re-introduction of work/school activities.
2	Light aerobic activity	Walking or stationary cycling at slow to medium pace. No resistance training.	Increase heart rate.
3	Sport-specific Exercise	Running or skating drills. No head impact activities.	Add movement.
4	Non-contact training drills	Harder training drills, e.g. passing drills. May start progressive resistance training.	Exercise, coordination and increased thinking.
5	Full contact Practice	Following medical clearance and complete return to school.	Restore confidence and assess functional skills by coaching staff.
6	Return to sport	Normal game play.	



The following is an outline of the *Return-to-Sport Strategy* that should be used to help athletes, coaches, trainers, and medical professionals to partner in allowing the athlete to make a gradual return to sport activities. Activities should be tailored to create a sport-specific strategy that helps the athlete return to their respective sport.

An initial period of 24-48 hours of rest is recommended before starting their *Sport-Specific Return-to-Sport Strategy*. If the athlete experiences new symptoms or worsening symptoms at any stage, they should go back to the previous stage. It is important that youth and adult student-athletes return to full-time school activities before progressing to stage 5 and 6 of the *Sport-Specific Return-to-Sport Strategy*. It is also important that all athletes provide their coach with a *Medical Clearance Letter* prior to returning to full contact sport activities.

1Source: McCrory et al. (2017). Consensus statement on concussion in sport – the 5th international conference on concussion in sport held in Berlin, October 2016. *British Journal of Sports Medicine*, *51*(11), 838-847. http://dx.doi.org/10.1136/bjsports-2017-